

Solubility and thermodynamic function of carbamazepine-saccharin (CBZ-SAC) co-crystal in ethanolic solution

Fatinah Ab Rahman ¹, Engku Nadia Engku Mat Nasir ¹, Syarifah Abd Rahim ^{1 *}, Raihana Zahirah Edros ², Nornizar Anuar ³

¹ *Faculty of Chemical & Natural Resources Engineering, Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300 Kuantan, Pahang, Malaysia*

² *Faculty of Engineering Technology, Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300 Kuantan, Pahang, Malaysia*

³ *Faculty of Chemical Engineering, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia*

Abstract:

The aim of study was to assess solubility and thermodynamic function of CBZ-SAC co-crystal in varying SAC concentrations in ethanolic solution at temperatures (25-50 oC). Experimental solubilities of CBZ-SAC co-crystal were determined by gravimetric and high performance liquid chromatography (HPLC) methods. Experimental solubilities of co-crystal in mole fraction from HPLC method were correlated well with Van't Hoff model. The solubilities of co-crystal were obtained highest in CBZ : SAC with ratio of 1.00 : 1.00 and the co-crystal solubility is higher than the pure form of CBZ (Fig.1) [1]. Based on the apparent thermodynamic analysis parameter, it was indicated an "endothermic and entropy-driven dissolution" of CBZ-SAC co-crystal [2,3,4]. From these results, CBZ-SAC co-crystal has been considered freely soluble in ethanol solution with non-SAC excess.